

Reflection on event horizon for collapsing black holes

M.Yu.Kuchiev*

School of Physics, University of New South Wales, Sydney 2052, Australia

(Dated: February 7, 2008)

It was argued recently that there exists an unexpected phenomenon, the reflection of incoming particles on the event horizon of black holes (Kuchiev(2003)). This means that a particle approaching the black hole can bounce back into the outside world due to those events that take part strictly on the horizon. Previously the effect was discussed in relation to eternal black holes. The present work shows that the effect exists for collapsing black holes as well.

PACS numbers: 04.70.Dy, 04.20.Gz

I. INTRODUCTION

An interest in quantum properties of the scattering problem that describes an impact of a particle on a black hole was inspired by the discovery of the Penrose process [1], and the works of Zel'dovich [2] and Misner [3] devoted to the energy extraction from the Kerr black hole. First numerical results for scattering of scalar particles on rotating black holes were presented by Press and Teukolsky [4, 5]. The analytical solution of the scattering problem, which had a strong influence on further developments, was found by Starobinsky [6] for the scalar field and Starobinsky and Churilov [7] for electromagnetic and gravitational waves scattered by the rotating Kerr black hole. Independently, Unruh [8] considered scattering of scalar and fermion particles on Schwarzschild black holes. Further study of the scattering problem was presented by Sanchez [9]. After these and the numerous works that followed, it has been assumed that the scattering problem is completely understood, see details and bibliography in the books [10, 11, 12, 13].

The present work discusses particular aspects of the scattering problem related to quantum phenomena, which take place strictly on the event horizon of black holes. The important relevant effects in this area are the Hawking radiation [14, 15], and the Unruh process [16]. The Hawking radiation is associated with the entropy of black holes, which, according to Bekenstein [17, 18, 19], equals the area of the horizon. For a review of the Hawking radiation and the thermodynamics properties of black holes see Refs. [20] and [21]. The Hawking radiation can be treated as a tunneling process, see e.g. Parikh and Wilczek [22], Khriplovich [23], and Khriplovich and Korokin [24] and references therein. Another line of research on quantum phenomena aims at quantization of energy levels of black holes, see discussion in Bekenstein [25, 26] and references therein. Treatment of a number of different quantum phenomena on the event horizon can be found in the review by 't Hooft [27] which, among other issues discusses the brick wall model of Ref. [28], aimed

at explanation of the entropy of black holes.

The effect of the Hawking radiation can be considered in terms of the influence that the horizon exercises on the vacuum of the theory. It has been realized recently that quantum events on the horizon have also a strong impact on the wave function of a particle that approaches a black hole [29, 30, 31]. As a result there arises a new qualitative feature in the scattering problem. A particle approaching the black hole can bounce on its event horizon back into the outside world. This phenomenon, which is referred to below as the reflection on the horizon (RH) using the terminology of [30], is due to pure quantum reasons; in the classical approximation it is obviously absent. The RH effect, which is strong when the wavelength of the particle exceeds the radius of the event horizon, reduces the absorption cross section, making it zero in the infrared region, as was demonstrated explicitly in Ref.[32] for scattering of low-energy scalar massless particles on Schwarzschild black holes. This behavior of the cross section differs qualitatively from the result of Unruh [8], which stated that this cross section equals the area of the horizon. Ref. [32] argued that similar reduction of the cross section is expected for scattering of any massless particle by any black hole in the low energy limit. Recent Ref. [33] gives a brief summary of progress related to the RH.

Refs.[29, 30, 31, 32, 33] suggested several arguments in favor of the RH. However, all of them, one way or another, were related to eternal black holes, inspiring a question as to whether the phenomenon exists for collapsing black holes as well, or is merely a peculiar feature relevant to eternal black holes only. The present work shows that the RH does take place for collapsars being thus a common effect for black holes.

II. SYMMETRY AND ANALYTICAL CONTINUATION

Let us formulate briefly and in general terms the main idea of this work. Consider a quantum system characterized by a set of coordinates Q , which include the time variable; generally speaking this set may be infinite. Take some quantum state of this system, not necessarily sta-

*Email:kuchiev@newt.phys.unsw.edu.au

tionary, which is described by the wave function $\Psi(Q)$. Assume now that there is some discrete symmetry that characterizes the system, calling the operator that generates this symmetry $\hat{S}$. It suffices to restrict out presentation to the simplest case, when the symmetry group is Z_2 , i.e. the symmetry group is generated by the only operator $\hat{S}$, which satisfies $\hat{S}^2 = 1$. (Though more general symmetries can also be covered by the method discussed, they will not play a role in the present work.) Since $\hat{S}$ is a symmetry, the function $\Psi'(Q) = \hat{S}[\Psi](Q)$ necessarily describes some physically allowed state of the system.

Let us consider now the complex extensions of the coordinates Q , using the analytical continuation for constructing the wave function $\Psi(Q)$ for the complex-valued coordinates. Take some closed contour C that runs in the multidimensional space of the complex coordinates Q . Fig. 1 gives a schematic presentation for this contour. The contour starts and finishes at some real physical point Q . Generically, the wave function $\Psi(Q)$ should have a sufficiently complex Riemann surface that possesses some singularities; the cuts attached to them give access to different sheets of the Riemann surface. Correspondingly, the value of the wave function depends on the sheet of the Riemann surface where it is taken. Thus, if the contour C crosses some cut that separates two sheets of the Riemann surface, called A_1 and A_2 in Fig. 1, then after returning to the initial physical value of Q the wave function acquires a new value, call it $\tilde{\Psi}(Q)$. Assume now




FIG. 1: Schematic presentation of the analytical continuation of the wave function $\Psi(Q)$ into the complex-valued coordinates Q . The contour C goes from the sheet A_1 of the Riemann surface over the cut to another sheet A_2 . As a result the wave function changes from $\Psi(Q)$ into $\tilde{\Psi}(Q)$.

that one is able to verify that the analytical continuation along the contour C and the symmetry transformation with the operator $\hat{S}$ have one and the same influence on the system. Then one can say that

$$\hat{S}[\Psi](Q) = \tilde{\Psi}(Q) . \quad (2.1)$$

This equality is derived from the initial wave function $\Psi(Q)$ by the symmetry transformation, which is implemented on the left-hand side directly, while on the right-hand side it works through the analytical continuation. One concludes from this that Eq.(2.1) provides a symmetry condition on $\Psi(Q)$. Eq.(2.1) is based on the relation between a monodromy transformation (analytical continuation) and a discrete symmetry of the system, which generally speaking is a well known method. To mention some result derived with its help one can remember the

known Seiberg-Witten solution for the $N = 2$ supersymmetric theory Ref.[34].

The system discussed in this work consists of collapsing matter and an incoming probing particle. The relevant discrete symmetry of the system is the time inversion, which transforms the collapsing matter (black hole) into an expanding matter (white hole), and at the same time forces the incoming particle to become the outgoing one. We will see in Section VIII that Eq.(2.1) puts a restriction on the wave function of the particle, deriving from it that the RH really takes place.

III. COLLAPSING BLACK HOLE

This section summarizes briefly several important for us, though well known facts related to the collapse of the dust matter, i. e. matter without pressure, $p = 0$, see books [35, 36] for details. For simplicity let us take the spherically symmetrical case when the collapse can be described by the Tolman metric

$$ds^2 = -d\tau^2 + \exp(\lambda) dR^2 + r^2 d\Omega^2 , \quad (3.1)$$

which is a function of τ , the radial variable R , and angular variables θ, φ ; $d\Omega^2 = d\theta^2 + \sin^2 \theta d\varphi^2$. The functions $\lambda = \lambda(R, \tau)$ and $r = r(R, \tau)$ are defined by the Einstein equations, which give for them

$$\exp(\lambda) = \frac{r'^2}{f(R) + 1} , \quad (3.2)$$

$$\dot{r}^2 = f(R) + F(R)/r . \quad (3.3)$$

Here and below the dotted and primed functions indicate derivatives over τ and R . The two functions $f(R)$, $F(R)$ provide the initial conditions for the spherically symmetrical dust matter. One of them, namely $F(R)$, defines the density of matter ϵ

$$8\pi G\epsilon = \frac{F'}{r'^2} . \quad (3.4)$$

The total mass of matter inside a sphere with the radius R equals

$$m(R) = 4\pi \int_0^R \epsilon r^2 dr = \frac{F(R)}{2G} . \quad (3.5)$$

The integrand here has no additional factors of $\exp(\lambda)$ or r' due to the defect of mass caused by gravity. Eq.(3.5) shows that $F(R)$ can be considered as the gravitational radius for the mass of matter accumulated inside the sphere of radius R , $F(R) = 2Gm(R)$. Another function $f(R)$, which appears in the formulation of the problem, gives an additional parameter that is needed to define the distribution of velocities of the matter given by Eq.(3.3). The two functions $f(R)$, $F(R)$ will be presumed to satisfy conditions

$$f(R) + 1 > 0 , \quad F(R) > 0 , \quad (3.6)$$

$$r' \geq 0 , \quad F'(R) \geq 0 . \quad (3.7)$$

Eqs.(3.6) are necessary to make $\exp(\lambda)$ and r positive, whereas Eqs.(3.7) exclude crossings of different spherical shells of the collapsing matter, which simplifies discussion.

Each shell of matter is characterized by some value of R , which remains constant during the collapse

$$R = \text{const} . \quad (3.8)$$

Integrating Eq.(3.3) when Eq.(3.8) holds one finds the explicit relation that defines the radius vector $r = r(R, \tau)$ of the shell of matter

$$\tau = - \int_{r_0}^r \frac{dr}{(f + F/r)^{1/2}} . \quad (3.9)$$

Here $r_0 = r_0(R)$ is an arbitrary function of R . The sign minus in front of the integral in Eq.(3.9) indicates that all velocities of the collapsing matter are negative, $\dot{r} < 0$. A fixed value of R in Eq.(3.9) prompts a shortcut notation $f = f(R)$, $F = F(R)$, which is also used below. If one changes the sign in front of the integral in Eq.(3.9), then the metric in Eq.(3.1) describes the exploding matter i.e. the white hole, which plays a significant role below.

In the particular case, when

$$f(R) = 0 , \quad F(R) = r_g = \text{const} , \quad (3.10)$$

the Tolman metric Eq.(3.1) represents the metric of eternal black holes. To see this clearly, one can choose $r_0 = r_g[3R/(2r_g)]^{3/2}$, finding that $r = r_g[3/(2r_g)(R - \tau)]^{3/2}$. Then Eq.(3.1) reduces to the Lemaitre metric

$$ds^2 = -d\tau^2 + (r_g/r) dR^2 + r^2 d\Omega^2 . \quad (3.11)$$

Further transformation of variables $R, \tau \rightarrow r, t$

$$\tau = \pm t \pm \int \frac{(r_g/r)^{1/2}}{1 - r_g/r} dr, \quad R = r + \int \frac{(r/r_g)^{1/2}}{1 - r_g/r} dr, \quad (3.12)$$

converts the Lemaitre metric to the Schwarzschild metric

$$ds^2 = - \left(1 - \frac{r_g}{r}\right) dt^2 + \frac{dr^2}{1 - r_g/r} + r^2 d\Omega^2 , \quad (3.13)$$

which explicitly describes a static eternal black hole.

IV. COMPLEX TRANSFORMATIONS AND TIME INVERSION FOR COLLAPSING MATTER

Let us introduce a complex structure associated with the Tolman metric Eq.(3.1). The angular variables θ, φ would not play a role below because we restrict discussion to the spherically symmetrical case. Our aim is to generalize the radial motion, allowing the radial variables to acquire complex values. It is sufficient to keep R real, allowing only r and τ to become complex-valued. It suffices also to assume that the classical equations of motion for matter Eqs.(3.8),(3.9) hold; in other words, we consider here the analytical continuation of the geodesics for the collapsing matter.

Eq.(3.9) allows one to take one of the variables r, τ as an independent variable, and treat the other one as its function. The conventional approach (which was followed above) takes r as a function of τ . However, it is more convenient here to assume that τ is a function of r , $\tau = \tau(r)$, see Fig. 2. This choice allows one to keep connection with the case of eternal static black holes, where the time variable does not play a role and the analytical continuation is applied to r , see Section VII. Obviously the function $\tau(r)$ presented by the integral in




FIG. 2: The Riemann surface for the function $\tau(r)$ possesses the cut of the squareroot nature that connects the points $r = 0$ and $r = -F/f$. Similar cut exists for the action $S(r, \tau(r))$ of the probing particle. Additionally, the action has the logarithmic singularity on the horizon $r = F(R)$ with the corresponding cut attached. The analytical continuation along the contour C generates the time inversion Eq.(4.1) for the collapsing matter; additionally, it results in transformations Eqs.(6.5),(6.6) for the action of the probing particle.

Eq.(3.9) can be expressed via elementary functions, but it is more instructive to derive its analytical properties directly from Eq.(3.9). On the complex plane r the integral in Eq.(3.9) has two singular points, $r = 0$ and $r = -F/f$, which match singularities of the radical $(f + F/r)^{1/2}$ in the integrand. This makes $\tau(r)$ an analytical function of r , which has a cut on the complex plane r that runs between the two singular points $r = 0$ and $r = -F/f$. Fig. 2 shows this cut running over the section $(-F/f, 0)$ of the real axis. Using the analytical continuation over this cut one constructs the Riemann surface for this function. The singularities at $r = 0$ and $r = -F/f$ are of the squareroot nature. Therefore the Riemann surface includes two Riemann sheets.

Let us assume that the initial real physical values of r and τ belong to the outside region, i. e. $r \geq F(R)$, where $R = R(r, \tau)$. Starting from this point one can continue the function $\tau(r)$ along the contour C that crosses the cut and then returns to the initial physical value of r , as shown in Fig. 2. Under this transformation (usually called a monodromy) the function is transformed as

$$\tau(r) \rightarrow -\tau(r) + \alpha . \quad (4.1)$$

Here the sign minus arises due to the squareroot nature of the crossed cut. The term α in Eq.(4.1) is a constant, which depends on r_0 in Eq.(3.9). Since r_0 is real, α is

also real and can be absorbed in the shift of the origin of time.

Eq.(4.1) shows that the analytical continuation described results in the inversion of the time variable. The equations of motion of the collapsing matter under this continuation become equations of motion for the expanding matter. Speaking plainly, this analytical continuation transforms a black hole into a white hole.

V. PROBING PARTICLE

Consider motion of a probing particle in the metric Eq.(3.1) created by the collapsing matter. It suffices to restrict discussion to the simplest case of a pure radial motion of the massless scalar particle, i.e. to choose the spin s , orbital momentum l , and mass μ of the particle to be all zero, $s = l = \mu = 0$. The corresponding classical action is a function of R and τ only, $S = S(R, \tau)$. The Hamilton-Jacobi equations of motion $g^{\mu\nu} \partial_\mu S \partial_\nu S = 0$, which follow from Eq. (3.1) in this case, have a simple form

$$\frac{\partial S}{\partial \tau} = \pm \frac{1}{r'} (f + 1)^{1/2} \frac{\partial S}{\partial R}. \quad (5.1)$$

Here $f = f(R)$; r is defined in Eq.(3.9) as a function of R and τ , $r = r(R, \tau)$; the signs plus and minus correspond to the outgoing and incoming trajectories respectively. Eq.(5.1) is formulated in terms of the initial variables R, τ of the Tolman metric. It is convenient now to look at the problem from another perspective, considering r, τ as independent variables and assuming that $R = R(r, \tau)$ is a function defined by the geodesics Eq. (3.9). One easily finds that

$$\left(\frac{\partial S}{\partial R} \right)_\tau = r' \left(\frac{\partial S}{\partial r} \right)_\tau, \quad (5.2)$$

$$\left(\frac{\partial S}{\partial \tau} \right)_R = \left(\frac{\partial S}{\partial \tau} \right)_r - r' \dot{R} \left(\frac{\partial S}{\partial r} \right)_\tau, \quad (5.3)$$

where $\dot{R} = (\partial R / \partial \tau)_r$. Differentiating Eq.(3.9), while keeping either $\tau = \text{const}$, or $r = \text{const}$, one finds also that

$$r' \dot{R} = (f + F/r)^{1/2}. \quad (5.4)$$

Using Eqs.(5.2),(5.3) and (5.4) one rewrites Eq.(5.1) in a more convenient form

$$- \frac{\partial S}{\partial \tau} = w \frac{\partial S}{\partial r}, \quad (5.5)$$

where $S = S(r, \tau)$ and the function w is defined as

$$w \equiv w_\pm(r, \tau) = - \left((f + F/r)^{1/2} \mp (f + 1)^{1/2} \right). \quad (5.6)$$

The signs plus and minus here match the signs in Eq.(5.1), the variables $f = f(R)$ and $F = F(R)$ are both functions of r, τ since it is presumed that $R = R(r, \tau)$.

To clarify the physical meaning of the factor w in Eq.(5.5) it is convenient to use the canonical formalism introducing the Hamiltonian H and the momentum of the particle p

$$p = \frac{\partial S}{\partial r}, \quad (5.7)$$

$$H = - \frac{\partial S}{\partial \tau} = wp, \quad (5.8)$$

where the last identity follows from Eq.(5.5). Then from Eq.(5.8) and the canonical equations of motion one finds that w equals the velocity of the particle

$$\frac{dr}{d\tau} = \frac{\partial H}{\partial p} = w, \quad (5.9)$$

which makes Eq.(5.5) convenient for the following discussion.

VI. COMPLEX TRANSFORMATIONS FOR PROBING PARTICLE

Let us discuss an analytical continuation of the variables, which describe the radial motion with zero orbital momentum of the scalar massless probing particle. This task is similar to the one dealt with in Section IV, where the analytical continuation was applied to the collapsing matter. However the study of the probing particle will be more detailed. In Section IV we discussed the analytical continuation assuming that the equations of motion are satisfied (i.e. assuming that Eqs.(3.8),(3.9) are valid). For the probing particle discussed here the analytical continuation of the action of the probing particle is considered. This presumes that the classical equations of motion for this particle are not necessarily valid. The classical action presents a particular interest since it allows one to construct the semi-classical wave function, see Section VIII.

Since the variables r, τ are not presumed to satisfy the equations of motion, the action $S(r, \tau)$ that describes the radial motion is a function of two independent variables. An analytical structure associated with this function for complex values of both variables is too sophisticated for our purposes. It is desirable therefore to simplify the problem forcing the action to be a function of only one complex variable.

With this in mind let us adopt the following approach. First, take some arbitrary real physical values for r, τ , call them $r = r_0$ and $\tau = \tau_0$. Then find a real physical R that satisfies classical equations of motion for the collapsing matter Eqs.(3.8),(3.9). In other words, find a shell of matter labeled by R , which at the moment τ_0 is located at the radius-vector r_0 . This makes R a function of the chosen physical variables $R = R(r_0, \tau_0)$. After that let us fix the found value of R and allow r and τ to take only particular complex values that satisfy Eq.(3.9).

Thus, the described procedure presumes that the complex values of r, τ satisfy the classical equations of motion (3.9) for matter. The complex radius vector r of the particle follows the complex extension of the geodesics. Under this condition the complex-valued τ can be taken as a function of the complex-valued r , $\tau = \tau(r)$. At the same time, there is no restriction on real physical values of r_0, τ_0 ; they remain two independent physical variables. The procedure outlined ensures that during the analytical continuation the action remains a function of only one complex variable r , $S(r, \tau) = S(r, \tau(r))$, which simplifies its analytical properties.

There exist, obviously, a variety of other ways to put some restriction on complex values of r and τ making τ a function of r , and thus forcing the action to be a function of one variable. An advantage of the procedure adopted here is that it matches the motion of matter, which follows the geodesics. This fact will allow one to combine results for the analytical continuation of the motion of the matter and probing particle.

It becomes clear below that the horizon plays an important role in the analytical structure of the action. Anticipating this feature one can presume that the physical values of r_0, τ_0 are chosen close to the event horizon, i. e. the following condition is fulfilled

$$|r_0 - F(R)| \ll F(R), \quad (6.1)$$

where R is a function of r_0, τ_0 as described above. This means that at the moment τ_0 the shell marked by R is close to its horizon as well. The subscripts in r_0, τ_0 are suppressed below; it is to be clear from the context whether real physical values of the variables r, τ , or their complex extensions are discussed.

Eq.(5.6) defines w as a function of r, τ . Following the procedure adopted, i. e. taking $R = \text{const}$, one makes w a function of r only. Then Eq.(5.6) shows explicitly that on the complex plane r this is a regular function with a squareroot-type cut from $r = 0$ to $r = -F/f$. This function takes the values w_+ and w_- on the two sheets of the corresponding Riemann surface. It is convenient to label these sheets as A_+ and A_- , and call them the sheet of outgoing particles, and the sheet of incoming particles respectively (keeping in mind that the velocities satisfy $w_+ > 0, w_- < 0$).

Since the action $S(r, \tau)$ is a solution of Eq.(5.5), which is governed by w , it necessarily exhibits properties similar to w . This means that it is an analytical function, which possesses the cut of the squareroot nature from $r = 0$ to $r = -F/f$ on the complex plane r , taking the values $S_+(r, \tau(r))$ and $S_-(r, \tau(r))$ on the two sheets of the Riemann surface. Here $S_+(r, \tau)$ is defined on the sheet A_+ and describes the outgoing motion, while $S_-(r, \tau)$ is located on A_- and is related to the incoming motion of the particle.

Let us discuss now the second important feature of the action, which is related to the node of the function w . Eq.(5.6) shows that this node is located on the horizon $r = F$ on the sheet A_+ of the Riemann surface, which

corresponds to the outgoing motion with $w = w_+$. This singularity of Eq.(5.5) enforces singular behavior on the action. The derivative $-\partial S/\partial \tau = H$ on the left-hand side of Eq. (5.5) represents the Hamiltonian, which is certainly positive $H > 0$ for all physical values of r, τ in the outside region $r > F$. It remains therefore nonnegative on the horizon $H \rightarrow H_0 \geq 0, r \rightarrow F$. One concludes that the derivative $\partial S/\partial r$ on the right-hand side of Eq. (5.5) exhibits a pole-type behavior on the horizon

$$\frac{\partial S}{\partial r} = \frac{H_0}{w} \rightarrow \frac{a}{r - F} + \text{const}, \quad \text{when } r \rightarrow F, \quad (6.2)$$

Here the residue is

$$a = 2(f + 1)^{1/2} F H_0 \geq 0, \quad (6.3)$$

as it follows from Eq.(5.6). The factor $H_0 = -(\partial S/\partial \tau)_{r=F}$ in Eq.(6.2) denotes the Hamiltonian of the particle on the horizon. From Eq.(6.2) one finds that the action itself has a logarithmic singularity on the horizon

$$S \rightarrow a \ln(r - F) + \text{const}, \quad r \rightarrow F. \quad (6.4)$$

We see that the analytical structure of the function $S(r, \tau(r))$ on the complex plane r is defined by two features shown in Fig. 2. There is a cut from $r = 0$ to $r = -F/f$ that connects the two sheets A_+ and A_- of the Riemann surface where the action takes the values $S_+(r, \tau)$ and $S_-(r, \tau)$ respectively. Additionally, there is the logarithmic singularity Eq.(6.4), which is located on the horizon being present on the sheet A_+ [41].

Consider now the transformation of the action when it is continued along the contour C in Fig. 2. The reasons inspiring to choose this contour for the analytical continuation are discussed below. To be specific let us take the action that describes the incoming trajectory $S_- = S_-(r, \tau(r))$. Correspondingly the contour C starts on the sheet A_- of the Riemann surface that corresponds to the incoming particles and, after crossing the cut over $(-F/f, 0)$, ends up on the sheet A_+ of the outgoing particles.

Combining the transformations arising from both the cut on the segment $(-F/f, 0)$ and the logarithmic singular point $r = F$, see Fig. 2, one concludes that the analytical continuation over the contour C results in the following transformation of the action

$$S_-(r, \tau) \rightarrow -S_+(r, \tau) + i\pi a. \quad (6.5)$$

Two features are to be noted here. First, the negative sign in front of $S_+(r, \tau)$, which is related to the crossing of the cut; the squareroot nature of this cut makes this sign inevitable. Second, there is the imaginary part in the right-hand side, which arises due to circling around the logarithmic singular point over the angle π counter clockwise. It adds $i\pi$ to the logarithmic function, forcing through Eq.(6.4) the imaginary constant $i\pi a$ to appear in the transformation of the action. Generally speaking, there may exist also an additional real additive constant

in the right-hand side of Eq.(6.5), but its calculation is not pursued here, since it is not important for our purposes. In contrast, the imaginary term plays a prominent role below.

Similarly the analytical continuation over the contour C for the action $S_+(r, \tau)$ results in the transformation

$$S_+(r, \tau) \rightarrow -S_-(r, \tau) - i\pi a. \quad (6.6)$$

The negative sign in front of the imaginary part is due to the fact that the logarithmic singularity in this example is circled in the clockwise direction.

In Section IV the analytical continuation that transforms the equations of motion for collapsing matter into equations of motion of the expanding matter was found. The results of the present Section are in line with this finding. The actions for incoming and outgoing particles transform one into another in Eqs.(6.5),(6.6).

VII. PARTICLE IN THE FIELD OF ETERNAL BLACK HOLE

It is instructive to specify the approach of Section VI for the simplest possible case, when both functions f, F , which govern the collapse are constants

$$f = \text{const}, \quad F = r_g = \text{const}. \quad (7.1)$$

This condition, which incorporates the physically interesting case of the eternal Schwarzschild black holes Eqs.(3.10),(3.13), indicates that w defined by Eq.(5.6) is a function of only one variable r , which allows to integrate Eq.(5.5) explicitly

$$S_{\pm}(r, \tau) = - \int^r \frac{\varepsilon dr}{(f + F/r)^{1/2} \mp (f + 1)^{1/2}} - \varepsilon \tau. \quad (7.2)$$

Here ε is the energy of the particle. The sign minus in front of the integral in Eq.(7.2) is related to negative velocities of the collapsing matter $\dot{r} < 0$, compare Eq.(3.9).

The explicit integral representation Eq.(7.2) allows one to verify that the action possesses the analytical structure shown in Fig. 2. Namely, on the complex plane r there exists a cut from $r = -F/f$ to $r = 0$, which connects the two sheets A_+ and A_- of the Riemann surface. On the sheet A_+ there exists an additional logarithmic singularity. It arises from the node in the integrand in Eq.(7.2), being located on the horizon

$$r = F \equiv r_g. \quad (7.3)$$

These facts confirm what is known from a more general approach of Section VI. One also re-derives Eqs.(6.5),(6.6) specifying that for eternal black holes a in Eq.(6.3) is a constant

$$a = 2\pi r_g \varepsilon = \frac{\varepsilon}{2T_H}, \quad (7.4)$$

where

$$T_H = \frac{1}{4\pi r_g} \quad (7.5)$$

is the Hawking temperature.

VIII. WAVE FUNCTION

Consider the scalar massless particle that moves with the zero orbital moment in the gravitational field of a collapsar, describing its propagation in the semiclassical approximation with the help of the wave function

$$\psi(r, t) = C \left(\exp(iS_-(r, t)) + \mathcal{R} \exp(iS_+(r, t)) \right). \quad (8.1)$$

Here $S_{\mp}(r, t)$ are the two actions that are related to the incoming and outgoing trajectories. Correspondingly, the first term in Eq.(8.1) describes the incoming wave, while the second one represents the outgoing wave.

The wave function Eq.(8.1) can be used to describe the process of absorption of particles by the black hole. In this case one needs to consider the behavior of this wave function on the horizon of each shell, i.e. to presume that r, τ satisfy condition

$$|r - F(R)| \ll F(R), \quad (8.2)$$

(which is similar to Eq.(6.1)). Eq.(8.2) has a clear physical meaning. It states that there is a shell of matter characterized by some particular value of $R = R(r, \tau)$, which at the moment of time τ has the radius vector r that is close to the horizon of this shell $F(R)$. Same can be rephrased as a statement that the amount of dust matter that is accumulated inside the sphere of radius r at the moment τ is close to a critical value, which is necessary to create a black hole. Generically, as the time passes on each shell converges to its own horizon $F(R)$ (eventually crossing it). Therefore there exists a set of critical values r_c of the radius vector that are located in an interval $0 < r_c \leq F_{\max}$, $F_{\max} = \lim_{R \rightarrow \infty} F(R)$; if r is close to any such critical value, then one can always choose τ to satisfy Eq.(8.2). For static black holes the set of critical radii degenerates to the only value of the Schwarzschild radius $r_c = r_g$.

Conventional wisdom prompts one to assume that in the vicinity of the horizon there should be no reflected wave; whatever comes close to the horizon must be absorbed, simply because we deal with a black hole, the object that is presumed to be *absolutely* black. Thus, one could anticipate that the second term in Eq.(8.1) is absent, i.e. the factor $\mathcal{R}$, which we will call the reflection coefficient, is zero, $\mathcal{R} = 0$ [42]. The absence of the reflected wave on the horizon, often called the Matzner condition in the stationary problems, see the references in the book [13], played a fundamental role in the scattering problem; it was used in the pioneering Refs. [4, 5, 6, 7, 8] as well as in numerous subsequent developments related to the scattering problem.

It turns out, however, that the reflection coefficient proves be nonzero $|\mathcal{R}| > 0$, the second term is present in Eq.(8.1) on the horizon. In other words there is the RH, as was shown in [29, 30, 31] for eternal black holes.

In order to verify this claim for the collapsing case let us apply the method outlined in Section II, which culminates in Eq.(2.1) for the wave function $\Psi(Q)$ of the quantum system. The system considered here includes the collapsing dust matter plus the incoming probing particle. However, quantum treatment is necessary only for the particle, as the dust matter is much less affected by quantum effects. This happens because the quantum effects prove strong only for low energies of a particle that approaches the horizon (see Eq.(8.7) below), while each drop of the collapsing dust can be considered as a heavy particle that has high energy associated with its mass. Anticipating this fact from the very beginning, one can describe the matter in the pure classical approximation, taking quantum effects into account only in the wave function of the particle $\psi(r, t)$.

Thus, Eq.(2.1) needs to be reformulated in terms of the wave function $\psi(r, t)$ of the particle, which moves in the given time-dependent metric produced by the collapsing matter. As the first step in this direction, let us use the contour C shown in Fig. 2 for the analytical continuation. Firstly, using this contour one continues analytically the equations of motion Eqs.(3.8),(3.9) for each shell R of the dust matter, converting the equations for the collapsing matter into equations of the expanding matter (converting the black hole into the white hole). The way to do this was described in detail in Section IV. Secondly, using same contour C one can fulfill the necessary transformation of the wave function of the particle. Using Eqs.(6.5),(6.6) for the classical action one finds that this operation can be written as a transformation of the wave function $\psi(r, t) \rightarrow \tilde{\psi}(r, t)$, where

$$\tilde{\psi}(r, t) = C e^{i\gamma} \left(\rho \exp(-iS_-(r, t)) + \frac{\mathcal{R}}{\rho} e^{i\delta} \exp(-iS_+(r, t)) \right). \quad (8.3)$$

Here ρ is governed by the imaginary constant in the transformation of the action (6.5),(6.6)

$$\rho = \exp(-\pi a), \quad (8.4)$$

where a is defined in Eq.(6.3), while the phases γ, δ are related to the real constants in the transformation of the action, which were omitted in (6.5),(6.6). Clearly, the first main term in Eq.(8.3) describes the outgoing wave, indicating that this wave function describes the creation of the particle. This is a sensible result since the analytical continuation transforms the matter into the white hole, as discussed above. The white hole is able to create particles. Thus, the considered analytical continuation leads to the time inversion.

Since the time invariance is valid, one is certainly free to fulfill the inversion of time directly in the equations of

motion. By applying the inversion of time to the classical equations of motion for the dust matter one brings them back to the collapsing case. The inversion of time leads also to the complex conjugation of the wave function of the particle, i.e. to the transformation

$$\psi(r, t) \rightarrow \psi'(r, t) = \psi^*(r, t). \quad (8.5)$$

Thus the inversion of time can be fulfilled in two different ways. either by the analytical continuation, resulting in Eq. (8.3), or it can be done directly in the equations of motion, which leads to Eq.(8.5). Since these two ways should give the same physical result, the wave functions in Eqs.(8.3),(8.5) should be the same, or at worst differ by some phase factor. One derives from this

$$\tilde{\psi}(r, t) = e^{i\chi} \psi'(r, t) = e^{i\chi} \psi^*(r, t). \quad (8.6)$$

This equality presents a restriction on the wave function of the particle, which stems from the symmetry related to the time inversion (compare Eq.(2.1), which was written in general terms). From definitions of the wave functions Eqs. (8.1),(8.3) and Eq.(8.6) one derives that the reflection coefficient is necessarily nonzero

$$|\mathcal{R}| = \rho = \exp(-\pi a) > 0. \quad (8.7)$$

Thus, the invariance under the time inversion Eq.(8.6) makes the process of the RH inevitable, the horizon is able to reflect the incoming particles.

Derivation of Eq.(8.6) was based on a particular contour on the complex plane r shown in Fig. 2. This contour must necessarily cross the cut that connects the sheets A_+ and A_- of the Riemann surface, thus leading to the time inversion in the equations of motion of the matter Eq.(4.1) and transformation of the wave function Eq.(8.3). The chosen way of curling around the logarithmic singularity leads to a sensible result, $|\mathcal{R}| < 1$ in Eq.(8.7), while the opposite direction of circling around this singularity would result in the violation of unitarity $|\mathcal{R}| > 1$. Such dependence of the result on the contour is common in semiclassical problems, see the book [37].

Eq.(8.7) shows that the RH really takes place, the horizon reflects particles.

IX. DISCUSSION

A. Main result

Our main result is presented in Eqs.(8.1),(8.7), which show that the wave function of the particle approaching the black hole horizon has a component that describes the reflected wave. In other words, the particle can bounce on the horizon back into the outside world. According to Eq.(8.7) the probability of this effect equals

$$P = |\mathcal{R}|^2 = \exp(-2\pi a) > 0, \quad (9.1)$$

where the parameter a is expressed in Eq.(6.3) via the functions $f = f(R)$, $F = F(R)$, which define the collapse. Their argument $R = R(r, \tau)$ depends on the coordinates r, τ of the wave function. Additionally, a depends on the Hamiltonian of the particle $H_0 = -\partial S/\partial \tau$, which is taken on the horizon. The expression for a simplifies in the limit of the static Schwarzschild black hole, $f(R) \rightarrow 0$, $F(R) \rightarrow r_g = \text{const}$ when Eq.(7.4) is valid. In this case Eq.(9.1) reduces to

$$P = \exp(-\varepsilon/T_H) > 0, \quad (9.2)$$

where T_H is the Hawking temperature Eq.(7.5). Eq.(9.2) coincides with the result of [29, 30, 31] derived exclusively for the static case. Eq.(9.2) shows that the reflection is strong for low energy particles $\varepsilon < T_H$. In other words, in the low energy limit the horizons of black holes act as mirrors.

B. Symmetry condition

It is interesting to compare the symmetry condition Eq.(8.6) with a similar condition found for the stationary case in [29, 30, 31]. The latter works focused on the wave function of the particle, leaving aside the effect of the discussed transformation on the collapsing matter. This simplified the consideration, but raised several qualitative questions. First, the symmetry, as it was derived in [29, 30, 31], was formulated in terms of regions I and III of the Kruskal plane [38] (definitions of regions follow [35]). However, one could argue that the collapsing black hole moves from region I into II, and does not come across region III. It was unclear therefore whether the proposed symmetry condition would hold for collapsing black holes. Second, there are known several ways to define the topological correspondence between regions I and III, see Refs. [39, 40]. Refs. [29, 30, 31] suggested to use a particular correspondence (the most obvious), postponing a question of whether the symmetry condition remains valid for all other possible ways of the topological correspondence.

The symmetry transformation in the present work is discussed from a broader perspective (in addition to the fact that a more general collapsing case is considered).

The inversion of the time is discussed in relation to both the wave function of the probing particle, and the motion of the matter. This point of view enables one to formulate simple physical answers for the mentioned problems. First, the inversion of the time variable not only brings the particle to region III, but it also transforms the collapsing matter into the expanding white hole, thus making region III relevant to the matter as well. Second, for any sensible topological correspondence between regions I and III the operation of the time inversion remains a symmetry of the space-time. This symmetry guarantees that the presented approach remains applicable for any topological identification between these regions.

C. Relation to Aharonov-Bohm effect

The RH can be considered as a manifestation of the logarithmic singularity, which the action exhibits on the event horizon. At the same time the classical trajectories do not come across this singularity (as one easily verifies using Fig. 2). Therefore the singularity has no influence on the classical equations of motion, being important only in the action. This property of the problem has a similarity with the Aharonov-Bohm effect, as was mentioned in [30, 31] in relation to the case of eternal black holes. Earlier a similarity that exists between physical events in the vicinity of the horizon and the Aharonov-Bohm effect was discussed in Ref.[22].

X. CONCLUSION

It is shown that the reflection on the horizon of black holes is a general phenomenon, which manifests itself for the collapsing case as well as for eternal black holes. One of interesting consequences of this fact is related to the information paradox. Since any incoming wave can bounce on the horizon back into the outside world, the horizon cannot destroy completely all the incoming information, at least part of it inevitably returns into the outside world.

This work was supported by the Australian Research Council. Discussions with V.V.Flambaum are appreciated.

-
- [1] R. Penrose, *Revista. Nuovo. Cim.* **1**, 252 (1969).
 - [2] Ya. B. Zel'dovich, *Zh. Eksp. Teor. Fiz. Pis'ma Red.* **14**, 270 (1970) [*JETP Lett.* **14**, 180 (1971)]; *Zh. Eksp. Teor. Fiz.* **62**, 2076 (1972) [*Sov.Phys.JETP* **35**, 1085 (1972)].
 - [3] C. W. Misner, *Bull. Am. Phys. Soc.* **17**, 472 (1972).
 - [4] W.H. Press, S. A. Teukolsky, *Nature (London)* **238**, 211 (1972);
 - [5] W.H. Press and S. A. Teukolsky, *Astrophys. J.* **141**, 443 (1974).
 - [6] A. A. Starobinsky, *Zh. Eksp. Teor. Fiz.* **64**, 48 (1973) [*Sov.Phys.JETP* **37**, 28 (1973)].
 - [7] A. A. Starobinsky and C. M. Churilov, *Zh. Eksp. Teor. Fiz.* **65**, 3 (1973) [*Sov.Phys.JETP* **38**, 1 (1974)].
 - [8] W. G. Unruh, *Phys. Rev. D* **14**, 3251 (1976); thesis Princeton Univ., 1971 (unpublished) (available University Microfilms, Ann Arbor, Mich.).
 - [9] N. Sanchez, *Phys. Rev. D* **18**, 1030 (1977).
 - [10] V. P. Frolov and I. D. Novikov, *Black hole physics: basic concepts and new developments* (1998) Dordrecht; Boston: Kluwer.

- [11] K. S. Thorne, *Black hole and time warps* (1994) New York: Norton.
- [12] S. Chandrasekhar, *The Mathematical Theory of Black Holes* (1993) New York: Oxford University Press.
- [13] J. A. H. Futterman, F. A. Handler, and R. A. Matzner, *Scattering from black holes* (1988) Cambridge; New York: Cambridge University Press.
- [14] S. W. Hawking, Nature (London) **248**, 30 (1974).
- [15] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975).
- [16] W.G.Unruh, Phys.Rev.D **14**, 870 (1977)
- [17] J. D. Bekenstein, Lett. Nuovo Cimento **4**, 737 (1972).
- [18] J. D. Bekenstein, Phys. Rev. D **7**, 2333 (1973).
- [19] J.D.Bekenstein, Phys. Rev. D **9**, 3292 (1974).
- [20] T.Jacobson *Introduction to quantum fields in curved spacetime and the Hawking effect*, gr-qc/0308048.
- [21] R. M. Wald, Living Rev. Rel. **4**, 6 (2001).
- [22] M.K.Parikh and F.Wilczek, Phys.Rev.Lett. **85**, 5042 (2000)
- [23] I.B.Khriplovich, *Radiation of charged particles by charged black hole*, gr-qc/9812060
- [24] I.B. Khriplovich, R.V. Korkin, J.Exp.Theor.Phys. **94** 453(2002) [Zh.Eksp.Teor.Fiz. **94** 531 (2002), in Russian]
- [25] J. D. Bekenstein, In *2001: A Spacetime Odyssey*, eds. M. J. Duff and J. T. Liu (2002) World Scientific Publishing, Singapore, pp. 21-31; hep-th/0107045).
- [26] J. D. Bekenstein, In *Advances in the Interplay between Quantum and Gravity Physics*, eds. P. G. Bergmann and V. de Sabbata (2002) Kluwer, Dordrecht, pp. 1-26; gr-qc/0107049).
- [27] G.'t Hooft, J. Mod. Phys. A **11**, 4623 (1996); gr-qc/9607022.
- [28] G.'t Hooft, Nucl. Phys. B **256** 727 (1984).
- [29] M. Yu. Kuchiev, *Reflection from black holes*, gr-qc/0310008.
- [30] M. Yu. Kuchiev, Europhys. Lett. **65**, 445 (2004).
- [31] M. Yu. Kuchiev, Phys. Rev. D **69**, 124031 (2004).
- [32] M. Yu. Kuchiev and V. V. Flambaum, Phys. Rev. D **70**, 044022 (2004).
- [33] M. Yu. Kuchiev and V. V. Flambaum, *Reflection on event horizon and escape of particles from confinement inside black holes*, gr-qc/0407077.
- [34] N. Seiberg and E. Witten, Nucl.Phys. B **431** 484 (1994); Nucl.Phys. B **426** 19 (1994); Erratum-ibid. B **430** 485 (1994).
- [35] C. W. Misner, K. S. Thorne, J. A. Wheeler, *Gravitation* (1973) San Fransisco: W. H. Freeman and Company.
- [36] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, 4th edition, (1975) Pergamon.
- [37] L. D. Landau and E. M. Lifshits, *Quantum mechanics: non-relativistic theory* (1977) Pergamon.
- [38] M. D. Kruskal, Phys. Rev. **119**, 1743 (1960).
- [39] G.W. Gibbons, Nucl. Phys. B **271**, 497 (1986).
- [40] A. Chamblin and G.W. Gibbons, Phys. Rev. D **55**, 2177 (1995); gr-qc/9607079.
- [41] The cut attached to this logarithmic singularity gives access to an infinite set of additional sheets of the Riemann surface, though they will not play a role in the following discussion.
- [42] Eq.(8.1) is written in the semiclassical approximation. In regions where the semiclassical approach fails the second term in the wave function is definitely present, describing the flux of particles scattered by those parts of the gravitational potential that violate the semiclassical conditions. In relation to the Hawking radiation this phenomenon is often referred to as the grey body effect; for low energy particles it takes place at large distances from a black hole. In contrast, in the close vicinity of the horizon one should expect the semiclassical method to work well (as can be easily verified for eternal black holes). The appearance of the second term here would indicate the reflection on the horizon.